

# Partial Meniscectomy

## Post Operative Instructions

James M. Bothwell, MD  
Fort Worth Orthopedics  
817.433.3450

### Medications

#### **\*Pain** (Tylenol with codeine, Tramadol, or Talwin)

You have been prescribed a strong narcotic analgesic for pain. Take regularly as prescribed and as needed for pain. A common side effect of this medicine is constipation. Please make sure you add fiber or a stool softener if necessary and drink plenty of fluids to help avoid this.

#### **\*Non-Steroidal Anti-Inflammatory** (Ibuprofen)

Unless otherwise directed, take Ibuprofen 600mg every 8 hrs with food for the first 7 days following surgery. ***Do not take this medicine if you are taking a blood thinner (Plavix, Coumadin), or have a history of an ulcer, bleeding disorder, kidney problems or drug allergy.***

#### **\*Antibiotic** (Keflex, Clindamycin)

If prescribed, please take this medicine as instructed.

#### **\*Antiemetic** (Phenergan)

This medicine should help if you are nauseated after your procedure. If you are unable to keep this medicine down please call the office as there may be an alternative.

### Diet

Keep to a light diet the day of surgery. You may advance to your usual diet the day after surgery.

### Activity

Keep your leg elevated. Try to keep your leg (knee and foot) above the level of your head as much as possible for at least the first 7 days after surgery. Be sure to move your ankles up and down (ankle pumps) to keep circulation moving in both your calves. *A good rule of thumb is a set of 20 every 20 minutes while awake.* You may move around on your crutches when you are comfortable. **Do not bear weight on your operative leg until your first post-op visit.** You may progress to weight bearing after your post-op visit. Your total recovery time is approximately 3-5 months. Further instructions regarding this will be given at your first postoperative visit.

## **Cold Therapy**

You may have been sent home with an Iceman for your knee. Use frequently for the first 10 days after surgery. Make sure pad does not directly touch skin.

*For questions and problems with the Iceman call Texas Brace System 817-416-7400, 24 hour on call line.*

## **Dressing Change**

Unless otherwise directed, your dressing will be changed at your first post op visit.

## **Showering**

Do not get the dressing or your incisions wet. You may use an occlusive dressing (plastic bag, plastic wrap, tape) until otherwise directed.

## **Physical Therapy**

Will likely start after your first post-op visit. Prescriptions and instructions will be given at that visit.

## **Post Operative Appointment:**

Your first post-op appointment will be 3-5 days after surgery.

## **Precautions**

Pain, bruising, tightness, and/or swelling in the knee are common symptoms to expect after surgery. However, **if you experience any signs of calf pain, please call our office immediately to speak with a provider.**

### To Do List:

- ✓ Ice for Iceman
- ✓ Garbage bags and tape for shower